

REMARKS ON POTENTIAL VERSUS BARRIER FUNCTION METHODS FOR LINEAR PROGRAMMING*

Kaoru Tone
Saitama University

(Received July 30, 1990)

Abstract We will point out some relations between potential and barrier function methods for linear programming. Then, based on the relations, we will show that the classical logarithmic barrier function method for linear programming can be adjusted so that it generates the optimal solution in $O(\sqrt{n}L)$ iterations, where n is the number of variables and L is the data length. The method can be seen as a barrier function version of Ye's "An $O(n^3L)$ potential reduction algorithm for linear programming".

1. Introduction

Since the epoch-making breakthrough by Karmarkar [12], the interior point methods for linear programming have been extensively studied in many aspects. One of the focuses of the studies is on the central trajectory leading to the optimal point. (See, for example, Sonnevend [19], Renegar [17], Bayer and Lagarias [1], Megiddo [15], Kojima, Mizuno and Yoshise [13], Monteiro and Adler [16], Goldfarb and Liu [7], Ye and Todd [22], Todd and Ye [20].)

The algorithm dealing with the central trajectory can be classified into two groups: one that follows the central trajectory directly and the other that minimizes a substitute function of the problem so that the successive points of iteration remain in the proximity of the central trajectory consequently. Among the latter approaches, some are called large-step algorithms in the sense that the step size of the movement in an iteration has no *a priori* bound but is determined by minimizing the substitute function on a line segment.

There are several types of such functions. We will deal with two of them. One is the classical logarithmic barrier functions originated by Frisch [4] and studied by Fiacco and McCormick [2] as applied to linear programming by many authors. (See Gill, Murray, Saunders, Tomlin and Wright [6], Gonzaga [8], Kojima, Mizuno and Yoshise [14] and Roos and Vial [18].) The other is the modern potential function introduced by Karmarkar [12], which has been studied and extended by many researchers. (See Gonzaga [10], Ye [21], Freund [3], Todd and Ye [20], among others.)

Recently, Roos and Vial [18] have proposed an $O(nL)$ iteration large-step logarithmic barrier function algorithms and Ye [21] has developed an $O(\sqrt{n}L)$ iteration potential reduction algorithm based on the primal-dual potential function. (Freund [3] and Gonzaga [10] have presented similar results.) The $O(\sqrt{n}L)$ iteration seems to be the best theoretical bound as of November 1989.

* This is a revised version of the paper, "An $O(\sqrt{n}L)$ Iteration Large-step Logarithmic Barrier Function Algorithm for Linear Programming" appeared in the Institute of Statistical Mathematics Cooperative Research Report 29 "Nonlinear Optimization-Modeling and Algorithm-", pp. 81-98, March, 1991.

The purpose of this paper is to point out some relations between the potential function and the logarithmic barrier function and to present a new $O(\sqrt{n}L)$ iteration large-step logarithmic barrier function algorithm based on the observations. Although Gonzaga [9] has presented an algorithm with the same polynomial bound in the same track, the formula for the control of the parameter is different from the present method. Gonzaga reduces it by a fixed rate when a centering condition comes to be satisfied, while our method reduces it adaptively from iteration to iteration.

2. Barrier Function and Potential Function

We will deal with the primal form of the linear programming problem:

$$\langle P \rangle \quad \min\{c^T x : Ax = b, \quad x \geq 0\} \quad (2.1)$$

where A is an (m, n) matrix, b and c are m - and n -dimensional vectors respectively, x is the variable n -dimensional vector to be determined optimally and the symbol T denotes the transpose.

The dual form of $\langle P \rangle$ is expressed as

$$\langle D \rangle \quad \max\{b^T y : s = c - A^T y \geq 0\} \quad (2.2)$$

where s and y are variable n - and m -dimensional vectors respectively.

For all x and y that are feasible for $\langle P \rangle$ and $\langle D \rangle$, we have

$$b^T y \leq z^{\text{op}} \leq c^T x \quad (2.3)$$

where z^{op} denotes the minimal (maximal) objective value of $\langle P \rangle$ ($\langle D \rangle$). As far as notations are concerned, e denotes the vector of all ones. The upper case letter (X) designates the diagonal matrix of the vector (x) in lower case.

For $\langle P \rangle$ and $\langle D \rangle$, we assume that

(1) the relative interior of the feasible regions of $\langle P \rangle$ and $\langle D \rangle$ is nonempty and we have an interior feasible solution x^0 and y^0 for $\langle P \rangle$ and $\langle D \rangle$ such that

$$Ax^0 = b, \quad x^0 > 0 \quad (2.4)$$

and

$$s^0 = c - A^T y^0 > 0, \quad (2.5)$$

(2) A has full row rank,

and

(3) the objective function value $c^T x$ is not a constant on the feasible region.

Associated with $\langle P \rangle$, we consider the logarithmic barrier function

$$f(x, \mu) = \frac{c^T x}{\mu} - \sum_{j=1}^n \ln(x_j) \quad (2.6)$$

where μ is a positive parameter.

The function f is strictly convex on the relative interior of the feasible region and achieves a minimum value at a unique point in it. In contrast to the classical barrier function $f(x, \mu)$, several authors have been studying extensively other types of functions motivated by Kar-markar [12]. ([8], [20], [21], [22]).

We will consider here two of them:

the primal potential function for an interior feasible x

$$f_P(x, \underline{z}) = \rho \ln(c^T x - \underline{z}) - \sum_{j=1}^n \ln(x_j) \quad (2.7)$$

and

the primal-dual potential function for an interior feasible pair (x, y, s)

$$f_{PD}(x, s) = \rho \ln(x^T s) - \sum_{j=1}^n \ln(x_j s_j) \quad (2.8)$$

where \underline{z} is a lower bound to z^{op} and ρ is a positive parameter.

For a pair of interior feasible primal-dual solution (x, y, s) , let $\underline{z} = b^T y$, then we have a relation between the primal and the primal-dual potential functions:

$$f_{PD}(x, s) = f_P(x, \underline{z}) - \sum_{j=1}^n \ln(s_j). \quad (2.9)$$

For an interior feasible x^0 and a positive parameter μ^0 , the projected Newton (ascent) direction associated with f is given by $X^0 p$, where

$$p = \frac{X^0 s}{\mu^0} - e \quad (2.10)$$

$$s = c - A^T y \quad (2.11)$$

and

$$y = (A(X^0)^2 A^T)^{-1} A X^0 (X^0 c - \mu^0 e). \quad (2.12)$$

For an interior feasible x^0 and a lower bound \underline{z}^0 to z^{op} , the projected gradient direction associated with f_P is given by $X^0 p_P$, where

$$p_P = \frac{\rho}{c^T x^0 - \underline{z}^0} X^0 s - e \quad (2.13)$$

$$s = c - A^T y \quad (2.14)$$

and

$$y = (A(X^0)^2 A^T)^{-1} A X^0 \left(X^0 c - \frac{c^T x^0 - \underline{z}^0}{\rho} e \right). \quad (2.15)$$

For the derivation of the above formulae, see Hertog and Roos [11]. It is evident that if we choose the parameter μ^0 as

$$\mu^0 = \frac{c^T x^0 - \underline{z}^0}{\rho} \quad (2.16)$$

then we have

$$p = p_P. \quad (2.17)$$

This fact is the basis on which our algorithm stands.

Now, we consider a line segment in the interior feasible region of $\langle P \rangle$ starting from an interior feasible x^0 with a direction d . The line segment is expressed in the parameter β as

$$x(\beta) = x^0 + \beta d \quad (2.18)$$

where $\beta \geq 0$, $Ad = 0$ and $x(\beta) > 0$.

On the line segment, we define a “gap function” between the barrier function and the primal potential function as

$$g(\beta) = f(x^0 + \beta d, \mu^0) - f_P(x^0 + \beta d, \underline{z}^0). \quad (2.19)$$

Then, we have the lemma:

[Lemma 1]

If $\mu^0 = \frac{c^T x^0 - \underline{z}^0}{\rho}$, then $g(\beta)$ is increasing for $\beta (0 \leq \beta < \beta_\infty)$, where $\beta_\infty = \min\{-x_j^0/d_j : d_j < 0, j = 1, 2, \dots, n\}$.

Proof.

$$\frac{dg(\beta)}{d\beta} = \frac{\rho\beta(c^T d)^2}{(c^T x^0 - \underline{z}^0)(c^T(x^0 + \beta d) - \underline{z}^0)} \geq 0.$$

Q.E.D.

Moreover, it is easy to see that $g(\beta)$ is convex in β .

Lemma 1 means the following:

[Lemma 2]

If $\mu^0 = \frac{c^T x^0 - \underline{z}^0}{\rho}$ and the logarithmic barrier function $f(x^0 + \beta d, \mu^0)$ decreases a certain amount from $f(x^0, \mu^0)$ on the line segment $x^0 + \beta d$ at $\beta = \beta^0$, then more reduction can be expected in the primal potential function $f_P(x^0 + \beta d, \underline{z}^0)$ from $f_P(x^0, \underline{z}^0)$ at $\beta = \beta^0$.

Based on the relations demonstrated by Lemma 2 and other lemmas explained later, we can construct an $O(\sqrt{n}L)$ iteration large-step logarithmic barrier function method which can be seen as a natural extension of Roos and Vial [18] and Ye [21].

3. Algorithm and Complexity

The following is a barrier function version of Ye's primal-dual algorithm [21] where the primal- or dual-step is chosen according to the 2-norm of pp . This algorithm generates successive pairs of interior feasible solutions $(x^0, s^0), (x^1, s^1), \dots$, from a given initial pair (x^0, s^0) . Since (x^{k+1}, s^{k+1}) is completely determined by (x^k, s^k) , we describe the algorithm as the process for generating (x^1, s^1) from (x^0, s^0) .

[Algorithm A]

Set $\rho = n + \nu\sqrt{n}$ with a constant $\nu \geq 1$ and

$$\alpha = 0.4. \quad (3.1)$$

Given x^0, s^0 and y^0 such that $Ax^0 = b, x^0 > 0$ and $s^0 = c - A^T y^0 > 0$,

$$(3.2)$$

Compute

$$\underline{z}^0 = b^T y^0 \quad (3.3)$$

$$\mu^0 = \frac{c^T x^0 - \underline{z}^0}{\rho} \quad (3.4)$$

$$y = (A(X^0)^2 A^T)^{-1} A X^0 (X^0 c - \mu^0 e). \quad (3.5)$$

$$s = c - A^T y \quad (3.6)$$

and

$$p = \frac{X^0 s}{\mu^0} \cdots e \quad (3.7)$$

If $\|p\| \geq \alpha$

then begin the primal-step as follows:

$$x^1 = x^0 - \beta^0 X^0 p \text{ with } \beta^0 = \operatorname{argmin}_{\beta \geq 0} f(x^0 - \beta X^0 p, \mu^0) \quad (3.8)$$

$$y^1 = y^0 \quad (3.9)$$

$$\underline{z}^1 = \underline{z}^0 \quad (3.10)$$

$$s^1 = s^0 \quad (3.11)$$

else begin the dual-step as follows:

$$x^1 = x^0 \quad (3.12)$$

$$y^1 = y \quad (3.13)$$

$$\underline{z}^1 = b^T y \quad (3.14)$$

$$s^1 = s \quad (3.15)$$

end.

The process terminates if the relation

$$c^T x^k - b^T y^k < 2^{-L} \quad (3.16)$$

is satisfied for some k .

The following two lemmas are essentially proved by Roos and Vial [18].

[Lemma 3]

If $\|p\| < \alpha$ then (y, s) , defined by (3.5) and (3.6), is an interior dual feasible solution and we have

$$c^T x^0 - z^{\text{op}} < c^T x^0 - b^T y \leq \mu^0 (n + \alpha \sqrt{n}). \quad (3.17)$$

Hence, noting $x^1 = x^0$, $\mu^0 = (c^T x^0 - \underline{z}^0)/(n + \nu \sqrt{n})$, $s^1 = s$ and $\underline{z}^1 = b^T y$, we have

$$(x^1)^T s^1 = c^T x^1 - \underline{z}^1 \leq \frac{n + \alpha \sqrt{n}}{n + \nu \sqrt{n}} (c^T x^0 - \underline{z}^0) = \frac{n + \alpha \sqrt{n}}{n + \nu \sqrt{n}} (x^0)^T s^0. \quad (3.18)$$

Thus, the duality gap is reduced at least by a factor $(n + \alpha \sqrt{n})/(n + \nu \sqrt{n}) (< 1)$.

Proof. See Appendix 1.

[Lemma 4]

If a step length of $\bar{\beta} = (1 + \|p\|_\infty)^{-1}$ is taken from x^0 along the direction $-X^0 p$, then the change in the barrier function f , denoted by Δf satisfies

$$\Delta f \leq -\|p\| + \ln(1 + \|p\|). \quad (3.19)$$

If $\|p\| \geq \alpha = 0.4$, then

$$\Delta f \leq -0.04. \quad (3.20)$$

Proof. See Roos and Vial [18].

The following two lemmas are derived from those by Ye [21].

[Lemma 5]

Let $0 < \gamma < 1/\sqrt{2}$ and $\rho = n + \nu\sqrt{n}$ with $\nu \geq 1$.

If $\|p\| < \gamma$, then we have

$$\begin{aligned} f_{PD}(x^0, s) &\leq f_{PD}(x^0, s^0) - \nu + \frac{\nu}{1+\nu}(\gamma+1) + \frac{\gamma^2(1+\gamma/\sqrt{1-\gamma^2})}{2(1-2\gamma^2)} \\ &\leq f_{PD}(x^0, s^0) - \frac{1}{2} + \frac{\gamma}{2} + \frac{\gamma^2(1+\gamma/\sqrt{1-\gamma^2})}{2(1-2\gamma^2)} \end{aligned} \quad (3.21)$$

where s is defined by (3.6).

If we set $\gamma = 0.4$, then the reduction in the primal-dual potential function is as follows:

$$f_{PD}(x^0, s) \leq f_{PD}(x^0, s^0) - 0.13. \quad (3.22)$$

Proof. See Appendix 2.

[Lemma 6]

Let $\rho = n + \nu\sqrt{n}$ with $\nu \geq 1$ and (x^k, s^k) ($k = 0, 1, 2, \dots$) be a series of interior primal-dual feasible solutions with $f_{PD}(x^0, s^0) = O(\sqrt{n}L)$. If, for a positive δ independent of n , the relation

$$f_{PD}(x^{k+1}, s^{k+1}) \leq f_{PD}(x^k, s^k) - \delta \quad (3.23)$$

holds for each k , then in $O(\nu\sqrt{n}L)$ iterations, we have

$$c^T x^k - b^T y^k = (x^k)^T s^k < 2^{-L}.$$

If $\nu = O(1)$ moreover, then the polynomial bound of iteration is $O(\sqrt{n}L)$.

Proof. See Appendix 3.

Now we are ready to show the theorem.

[Theorem 1]

If Algorithm A starts from an interior primal-dual feasible solution (x^0, s^0) with $f_{PD}(x^0, s^0) = O(\sqrt{n}L)$, then it terminates in $O(\sqrt{n}L)$ iterations.

Proof.

Let the series of the interior feasible primal-dual solutions generated by Algorithm A be (x^k, s^k) ($k = 0, 1, 2, \dots$). For each (x^k, s^k) , we have three potential functions f , f_P and f_{PD} defined by (2.6), (2.7) and (2.8) respectively. We will show that, for each iteration, the primal-dual potential function reduces at least by a positive value $\delta = 0.04$ and then we have the conclusion by Lemma 6.

The case $\|p\| \geq \alpha$:

In this case we move in the primal space from x^0 to x^1 as defined by (3.8). From Lemma 4, we have

$$f(x^1, \mu^0) - f(x^0, \mu^0) \leq -0.04.$$

Using the gap function g , the change in the primal potential function is expressed as

$$\begin{aligned} f_P(x^1, \underline{z}^0) - f_P(x^0, \underline{z}^0) &= f(x^1, \mu^0) - g(\beta^0) - f(x^0, \mu^0) + g(0) \\ &\leq -0.04 - g(\beta^0) + g(0). \end{aligned}$$

By Lemma 1, we have $g(\beta^0) \geq g(0)$.

Hence,

$$f_P(x^1, \underline{z}^0) - f_P(x^0, \underline{z}^0) \leq -0.04.$$

Noting $s^1 = s^0$ in this case, we have

$$\begin{aligned} f_{PD}(x^1, s^1) &= f_P(x^1, \underline{z}^0) - \sum_{j=1}^n \ln(s_j^0) \\ &\leq f_P(x^0, \underline{z}^0) - \sum_{j=1}^n \ln(s_j^0) - 0.04 \\ &\leq f_{PD}(x^0, s^0) - 0.04. \end{aligned}$$

The case $\|p\| < \alpha$:

From Lemma 5, we have

$$f_{PD}(x^1, s^1) \leq f_{PD}(x^0, s^0) - 0.13.$$

Q.E.D.

4. Concluding Remarks

We will point out several features of our algorithm.

4.1 On primal- and dual-step

In algorithm A, we choose either the primal or the dual step depending on $\|p\|$. Specifically, if $\|p\| \geq \alpha (= 0.4)$, then we employ the primal, otherwise the dual. The value 0.4 is not mandatory, but is used to assure a constant reduction in the primal-dual potential function even in the worst case. So, in the implementational phase of the algorithm, the following procedure may be recommendable:

If $\|p\| < 1$ and (x^0, s) (s defined by (3.6)) reduces f_{PD} a certain amount, then we go into the dual step, otherwise into the primal. Also, the minimization of f_{PD} with respect to \underline{z} may be considerable.

4.2 On updating \underline{z}

If $\|p\| < 1$, then we have, from (3.18),

$$z^{op} \geq \underline{z}^1 > \underline{z}^0. \quad (4.1)$$

Thus, we can update the lower bound strictly. This fact means that if we start from \underline{z}^0 that is very close to z^{op} , then the centering condition $\|p\| < 1$ rarely holds and so we have few chances to visit the dual step.

It should be noted that to be in a proximity of the center, as characterized by $\|p\| < 1$, is not the object or goal of the path-following algorithm, but just a stimulus. By choosing $\mu = (cx - \underline{z})/\rho$, we change the stimulus, in a sense, adaptively and continuously. This shows a sharp contrast to Roos and Vial [18] and Gonzaga [9] where the centering condition is a necessity to promote their “outer step”.

4.3 On the choice of p

Although we employ $\rho = n + \nu\sqrt{n}$ with $\nu \geq 1$, it is interesting to observe the case $\rho = \theta(n + \nu\sqrt{n})$, with $\theta > 1$. From Lemma 6, the polynomial bound of iteration is $O(nL)$, worse than $O(\sqrt{n}L)$ of the present algorithm. Then, if $\|p\| < \alpha$ and we go into the dual step, it holds

$$\begin{aligned} (x^1)^T s^1 &= c^T x^1 - \underline{z}^1 = c^T x^0 - \underline{z}^1 \leq \frac{n + \alpha\sqrt{n}}{\theta(n + \nu\sqrt{n})} (c^T x^0 - \underline{z}^0) < \frac{1}{\theta} (c^T x^0 - \underline{z}^0) \\ &= \frac{1}{\theta} (x^0)^T s^0. \end{aligned} \quad (4.2)$$

Thus, the duality gap reduces at least by a factor $1/\theta (< 1)$. If we set $\theta = 2$, then Algorithm A will behave similarly to Roos and Vial [18] although the correspondence is not exact.

4.4 On the step size of the algorithm

Algorithm A uses the logarithmic barrier function f to determine the step size in the primal step. If, instead, we employ the primal potential function f_P for this purpose, then Algorithm A coincides with Ye’s primal potential reduction algorithm [21]. In this context, it may be possible that other types of the substitute functions with the same polynomial bound exist.

As for the step size, let

$$\beta^0 = \operatorname{argmin}_{\beta \geq 0} f(x^0 - \beta X^0 p, \mu^0) \quad (4.3)$$

and

$$\beta^1 = \operatorname{argmin}_{\beta \geq 0} f_P(x^0 - \beta X^0 p, \underline{z}^0). \quad (4.4)$$

Then, we have

$$\beta^0 \leq \beta^1, \quad (4.5)$$

as otherwise Lemma 1 does not hold.

4.5 On the dual barrier function algorithm

The dual barrier function \bar{f} and the dual potential function f_D for linear programming are defined as

$$\bar{f}(y, \mu) = -\frac{b^T y}{\mu} - \sum_{j=1}^n \ln(s_j)$$

and

$$f_D(y, \bar{z}) = \rho \ln(\bar{z} - b^T y) - \sum_{j=1}^n \ln(s_j),$$

where μ and ρ are positive numbers, \bar{z} is an upper bound to the optimal objective value z^{op} and $s = c - A^T y$.

By considering the difference between \bar{f} and f_D in the same way as in the primal case, we can develop an $O(\sqrt{n}L)$ iteration large-step dual barrier function algorithm. See Appendix 4 in details. (cf. The dual algorithm in Ye [21].)

Acknowledgement

I wish to thank two referees for helpful remarks.

Appendix

Appendix 1. (Proof of Lemma 3.)(Roos and Vial [18])

Since

$$\|p\| = \left\| \frac{X^0 s}{\mu^0} - e \right\| < \alpha < 1,$$

(s, y) is an interior dual feasible solution and so $\underline{z}^1 = b^T y < z^{\text{op}}$.

On the other hand,

$$|e^T \left(\frac{X^0 s}{\mu^0} - e \right)| \leq \|e\| \left\| \frac{X^0 s}{\mu^0} - e \right\| \leq \alpha \sqrt{n}$$

and

$$e^T \left(\frac{X^0 s}{\mu^0} - e \right) = \frac{(x^0)^T s}{\mu^0} - n = \frac{c^T x^0 - b^T y}{\mu^0} - n.$$

Hence, noting $x^1 = x^0$, $\mu^0 = (c^T x^0 - \underline{z}^0)/(n + \nu\sqrt{n})$, $s^1 = s$ and $\underline{z}^1 = b^T y$, we have

$$(x^1)^T s^1 = c^T x^1 - \underline{z}^1 \leq \frac{n + \alpha\sqrt{n}}{n + \nu\sqrt{n}} (c^T x^0 - \underline{z}^0) = \frac{n + \alpha\sqrt{n}}{n + \nu\sqrt{n}} (x^0)^T s^0.$$

Q.E.D.

Appendix 2. (Proof of Lemma 5)

First, we will show a lemma.

[Lemma 7]

Given two numbers α and ν with $0 < \alpha < \nu$, the function

$$h(x) = x \ln((x + \alpha)/(x + \nu))$$

is decreasing for $x > 0$.

Proof.

$$h'(x) = \ln\left(1 + \frac{\alpha - \nu}{x + \nu}\right) + \frac{(\nu - \alpha)x}{(x + \alpha)(x + \nu)}.$$

Since $(\alpha - \nu)/(x + \nu) > -1$ for $x > 0$, we have

$$h'(x) \leq \frac{\alpha - \nu}{x + \nu} + \frac{(\nu - \alpha)x}{(x + \alpha)(x + \nu)} = \frac{\alpha(\alpha - \nu)}{(x + \alpha)(x + \nu)} < 0.$$

Q.E.D.

Now, we prove Lemma 5.

Ye [21, page 247] proved that if $\|p\| < \gamma\sqrt{n/(n+\gamma^2)}$ with $\gamma < 1$ then the following inequality holds

$$n \ln(x^0)^T s - \sum_{j=1}^n \ln(x_j^0 s_j) \leq n \ln(x^0)^T s^0 - \sum_{j=1}^n \ln(x_j^0 s_j^0) + \frac{\gamma^2}{2(1-\gamma)}.$$

Thus, for $\|p\| < \alpha$ with $\alpha < 1/\sqrt{2}$, we have

$$\begin{aligned} n \ln(x^0)^T s - \sum_{j=1}^n \ln(x_j^0 s_j) &\leq n \ln(x^0)^T s^0 - \sum_{j=1}^n \ln(x_j^0 s_j^0) + \frac{\alpha^2(1 + \alpha\sqrt{n/(n-\alpha^2)})}{2(1-\alpha^2-\alpha^2/n)} \\ &\leq n \ln(x^0)^T s^0 - \sum_{j=1}^n \ln(x_j^0 s_j^0) + \frac{\alpha^2(1 + \alpha/\sqrt{1-\alpha^2})}{2(1-2\alpha^2)}. \end{aligned} \quad (A1)$$

On the other hand, we have from (3.18),

$$\sqrt{n}(\ln(x^0)^T s - \ln(x^0)^T s^0) \leq \sqrt{n} \ln \left(\frac{n + \alpha\sqrt{n}}{n + \nu\sqrt{n}} \right). \quad (A2)$$

By Lemma 7 above, the right hand side of (A2) attains a maximum at $n = 1$ for $n \geq 1$. Thus,

$$\begin{aligned} \sqrt{n}(\ln(x^0)^T s - \ln(x^0)^T s^0) &\leq \ln \frac{1+\alpha}{1+\nu} = \ln(1 + \frac{\alpha-\nu}{1+\nu}) \\ &< \frac{\alpha-\nu}{1+\nu} = -1 + \frac{1+\alpha}{1+\nu}. \end{aligned} \quad (A3)$$

From (A1) and (A3), we have, for $\alpha < 1/\sqrt{2}$,

$$\begin{aligned} f_{PD}(x^0, s) &\leq f_{PD}(x^0, s^0) - \nu + \frac{\nu}{1+\nu}(\alpha+1) + \frac{\alpha^2(1 + \alpha/\sqrt{1-\alpha^2})}{2(1-2\alpha^2)} \\ &\leq f_{PD}(x^0, s^0) - \frac{1}{2} + \frac{\alpha}{2} + \frac{\alpha^2(1 + \alpha/\sqrt{1-\alpha^2})}{2(1-2\alpha^2)}, \end{aligned}$$

because $-\nu + \nu(\alpha+1)/(1+\nu)$ attains its maximum at $\nu = 1$ for $\nu \geq 1$.

Q.E.D.

Appendix 3. (Proof of Lemma 6)(Ye [21])

$$\begin{aligned} f_{PD}(x, s) &= \rho \ln(x^T s) - \sum_{j=1}^n \ln(x_j s_j) \\ &= (\rho - n) \ln(x^T s) - \sum_{j=1}^n \ln((x_j s_j)/(x^T s)). \end{aligned}$$

From the inequality of the geometric mean and the arithmetic mean, we have

$$-\sum_{j=1}^n \ln((x_j s_j)/(x^T s)) \geq n \ln n.$$

Hence,

$$\begin{aligned} (\rho - n) \ln(c^T x - b^T y) &= (\rho - n) \ln(x^T s) \leq f_{PD}(x, s) - n \ln n \\ &\leq f_{PD}(x, s). \end{aligned}$$

Thus, if we can reduce f_{PD} at least by δ (a constant independent of n), at each iteration, we have, after $(\rho - n)L/\delta$ iterations,

$$(\rho - n) \ln(c^T x - b^T y) \leq -(\rho - n)L + f_{PD}(x^0, s^0).$$

Assume that $f_{PD}(x^0, s^0) = O(\sqrt{n}L)$ and $\rho = n + \nu\sqrt{n}$, then after $O(\nu\sqrt{n}L)$ iterations, we have

$$c^T x - b^T y \leq 2^{-L}.$$

Q.E.D.

Appendix 4. The dual barrier function algorithm

[Algorithm B]

Set $\rho = n + \nu\sqrt{n}$ with a constant $\nu \geq 1$ and
 $\alpha = 0.4$.

Given x^0, s^0 and y^0 such that $Ax^0 = b$,

$x^0 > 0$ and $s^0 = c - A^T y^0 > 0$,

Compute

$$\bar{z}^0 = c^T x^0$$

$$\mu^0 = \frac{(x^0)^T s^0}{\rho}$$

$$x_1 = (S^0)^{-2} A^T (A(S^0)^{-2} A^T)^{-1} b$$

$$x_2 = (S^0)^{-1} (I - (S^0)^{-1} A^T (A(S^0)^{-2} A^T)^{-1} A(S^0)^{-1}) (S^0)^{-1} c$$

$$x = x_1 + \mu^0 x_2$$

and

$$p = -\frac{S^0 x}{\mu^0} + e.$$

If $\|p\| \geq \alpha$

then begin the dual-step as follows:

$$s^1 = s^0 + \beta^0 S^0 p$$

$$\text{with } \beta^0 = \operatorname{argmin}_{\beta \geq 0} \{(x^T S^0 p) \beta / \mu^0 - \sum_{j=1}^n \ln(1 + \beta p_j)\}$$

$$x^1 = x^0$$

else begin the primal-step as follows:

$$x^1 = x$$

$$s^1 = s^0$$

end.

The process terminates if $(x^k)^T s^k < 2^{-L}$ is satisfied for some k .

References

- [1] D. Bayer and J.C. Lagarias: The non-linear geometry of linear programming, Technical Reports, Bell Labs, Murray Hill, NJ, 1987.

- [2] A.V. Fiacco and G.P. McCormick: *Nonlinear Programming: Sequential Unconstrained Minimization Techniques*, John Wiley and Sons, New York, NY, 1968.
- [3] R.M. Freund: Polynomial-time algorithms for linear programming based only on primal scaling and projected gradients of a potential function, *Mathematical Programming*, Vol. 51 (1991), 203–222.
- [4] K.R. Frisch: The logarithmic potential method of convex programming, Technical Report, University Institute of Economics, Oslo, Norway, 1955.
- [5] G. de Ghellinck and J.-P. Vial: A polynomial Newton method for linear programming, *Algorithmica*, Vol. 1 (1986), 425–454.
- [6] P.E. Gill, W. Murray, M.A. Saunders, J.A. Tomlin and M.H. Wright: On projected Newton barrier methods for linear programming and an equivalence to Karmarkar's projective method, *Mathematical Programming*, Vol. 36 (1986), 183–209.
- [7] D. Goldfarb and S. Liu: An $O(n^3L)$ primal interior point algorithm for convex quadratic programming, Technical Report, Department of IEOR, Columbia University, New York, NY, 1988.
- [8] C. Gonzaga: An algorithm for solving linear programming problems in $O(n^3L)$ operations, In: N. Megiddo, ed., *Advances in Mathematical Programming-Interior Point and Related Methods*, Springer Verlag, 1989.
- [9] C. Gonzaga: Large-steps path-following methods for linear programming, Part I: barrier function method, *SIAM Journal on Optimization*, Vol. 1 (1991), 268–279.
- [10] C. Gonzaga: Large-steps path-following methods for linear programming, Part II: potential reduction method, *SIAM Journal on Optimization*, Vol. 1 (1991), 280–292.
- [11] D. den Hertog and C. Roos: A survey of search directions in interior point methods for linear programming, Report 89–65, Faculty of Technical Mathematics and Informatics, Delft University of Technology, The Netherlands, 1989.
- [12] N. Karmarkar: A new polynomial-time algorithm for linear programming, *Combinatorica*, Vol. 4 (1984), 373–395.
- [13] M. Kojima, S. Mizuno and A. Yoshise: A polynomial-time algorithm for a class of linear complementarity problems, *Mathematical Programming*, Vol. 44 (1989), 1–26.
- [14] M. Kojima, S. Mizuno and A. Yoshise: A primal-dual interior point algorithm for linear programming, In: N. Megiddo, ed., *Progress in Mathematical Programming — Interior Point and Related Methods*, Springer Verlag, 1989.
- [15] N. Megiddo: Pathways to the optimal set in linear programming, In: N. Megiddo, ed., *Progress in Mathematical Programming — Interior Point and Related Methods*, Springer Verlag, 1989.
- [16] R.C. Monteiro and I. Adler: Interior path following primal-dual algorithms. Part 1: Linear programming, *Mathematical Programming*, Vol. 44 (1989), 27–41.
- [17] J. Renegar: A polynomial-time algorithm, based on Newton's method, for linear programming, *Mathematical Programming*, Vol. 40 (1988), 59–93.
- [18] C. Roos and J.-Ph. Vial: Long steps with the logarithmic penalty barrier function in linear programming, Report 89–44, Faculty of Technical Mathematics and Informatics, Delft University of Technology, The Netherlands, 1989.
- [19] G. Sonnevend: An 'analytic center' for polyhedrons and new classes of global algorithms for linear (smooth, convex) programming, *Proc. 12th IFIP Conference on System Modeling and Optimization*, Budapest, 1985.
- [20] M.J. Todd and Y. Ye: A centered projective algorithm for linear programming, *Mathematics of Operations Research*, Vol. 15 (1990), 508–529.
- [21] Y. Ye: An $O(n^3L)$ potential reduction algorithm for linear programming, *Mathematical Programming*, Vol. 50 (1991), 239–258.

- [22] Y. Ye and M.J. Todd: Containing and shrinking ellipsoids in the path-following algorithm, *Mathematical Programming*, Vol. 47 (1990), 1–9.

Kaoru Tone
Graduate School of Policy Science
Saitama University
Urawa, Saitama 338
Japan